



Deterministic Entanglement of Trapped-Ion Spin-Qubits


Research supported by:
ARDA (P-43513-PH-QCO-02107-1)
E.U. (QGATES/CONQUEST)
EPSRC (QIP IRC)
Royal Society

J.P.Home, G.Imreh, B.C.Keith, D.M.Lucas, M.J.McDonnell, N.R.Thomas, D.N.Stacey, A.M.Steane and S.C.Webster
Centre for Quantum Computation, Clarendon Laboratory, Oxford University, U.K.




Main points

We present experiments and theory in quantum information processing using trapped ions.
This poster concentrates on entanglement and gates: see accompanying poster for cooling, coherence.




Summary of Results

- 10 two-ion (2 qubit) Rabi flops with high visibility
- Deterministic entanglement of 2 ions (calcium 40 spin qubits) at 82(2)% fidelity
- Schrodinger cat with 1 ion and motion:
 - up to $\langle n \rangle = 12$
 - well outside Lamb-Dicke regime: $\sigma_{2n} = 1$ preserved for 422 s with 80(20)% fidelity also $\sigma_{-2,0,+2}$ with 2 ions
- robust convenient tomography method
- (th.) factorization of general phase gates (ask for details)
- (th.) composite pulses for fast gate ($t = 1/\text{trap freq}$) insensitive to optical phase



Spin-dependent force


For two-qubit gates we use spin-dependent forces: push ions depending on spin state
Coulomb interaction gives a two-qubit phase.
The force is an optical dipole force in a standing wave with polarization gradient.



Difference frequency of laser beams oscillating force on ion

- A classical force displaces the motional state in phase space.
- In Lamb-Dicke regime, extent of motional wavepacket \ll the force appears to be spatially uniform, and an oscillating force drives the motional state around a circular loop in phase space. It returns to the origin after $t = 2/\Omega$, where Ω is the Rabi frequency.
- Outside the Lamb-Dicke regime, the trajectory of the motional state through phase space is modified, and the motional state is squeezed. The motional state returns to the origin early, i.e. $t_r < 2/\Omega$.


We simulate the NLD regime behaviour by numerical integration of Schrodinger's equation.



Schrödinger Cat experiments

Coherent states of a harmonic oscillator approximate to classical motion, and a superposition of such states at mesoscopic excitation $\langle n \rangle$ is a type of Schrödinger cat.


Oscillating spin-dependent force create such mesoscopic superpositions with single or pairs of ions. Spin state = measuring device entangled with the motion. We prove the 'cat' maintains its coherence by bringing the two parts back together and observing an interference. [As first demonstrated by Monroe *et al.* Science 272 1131]




Observe $P(\uparrow)$ as a function of ϕ .

$$P(\uparrow) = \frac{1}{2} (1 - \text{Re}[\langle \alpha_{\uparrow} | \alpha_{\downarrow} \rangle e^{i(\phi - \Delta\pi\tau)})]$$


where $\Delta\pi$ is a light shift due to off-resonant carrier excitation (due to σ component of the light field)



Fringe amplitude
 $A = |\langle \alpha_{\uparrow} | \alpha_{\downarrow} \rangle|$



Fringe phase
 $\text{Arg}(\langle \alpha_{\uparrow} | \alpha_{\downarrow} \rangle) + \Delta\pi\tau$



We observe cat states with $\langle n \rangle = 5.4$, $n_{\text{max}} = 4$, and a motional coherence time $T_2 = 170$ s.

Deterministic entanglement

- Deterministic** (i.e. single-shot, no post-selection) entanglement of 2 spin-qubits
- gate uses oscillating spin-dependent driving force used to create Schrödinger cats, with force frequency close to Ω_{str} & ion separation = integer number of standing wave periods
=> only stretch mode excited
=> states acquire a phase; do not.
- Gate operation


$$\hat{U} = \begin{pmatrix} 1 & & \\ & i & \\ & & i \\ & & & 1 \end{pmatrix}$$

(Leibfried *et al.* [Nature 422 412 (2003)].)


Experiments: $\Omega_{\text{com}} = 500$ kHz, ion sep = 9 nm = 22

Single pulse method: Implements gate + single qubit rotations (due to light shift).


For $\Omega_{\text{str}} = 2$ and $\Omega_{\text{com}} = 1/2$, we get the entangled state.



A further $\pi/2$ analysis pulse with variable phase demonstrates $\cos(2\phi)$ oscillations in the parity signal with amplitude > 0.5 .
Entangled state fidelity $> 75(5)\%$



Two pulse method: One pulse in each half of spin-echo. Single qubit rotations cancel. Entangled state density matrix reconstructed using tomography method (see below). Fidelity 82(2)%





Single-qubit gates, 1-2 ions

Spin qubit state coherently manipulated either by magnetic resonance or by stimulated Raman transition.

Single-ion Ramsey fringes

This data is for a two-pulse Ramsey sequence using magnetic resonance with a single trapped ion. Interference fringes are seen as the RF frequency is scanned.

Two-ion Rabi oscillations




2-ion spin-state = 2 qubits. Rabi flopping, here driven by the Raman transition (4.5 s time), gives a single-qubit rotation applied to both qubits simultaneously.

EIT readout infer $P(\uparrow\uparrow)$, $P(\uparrow\downarrow)$, $P(\downarrow\uparrow)$, $P(\downarrow\downarrow)$

Composite pulses for fast robust gates

Wobble gate works well at low Ω_{str} but is slow,
At high Ω_{str} both COM and STR modes excited, can't close both loops in a single pulse (incommensurate freq).
Tailor $f(t)$ in order to go faster? : lose insensitivity to optical phase


- We find fast composite pulse sequences maintaining insensitivity to ϕ_0
- Issues: loop closure, constant area, lightshift phase



$$f(t) = \sum_{n=1}^N T((t - t_n)/\tau_n) f_n \sin(\omega_n t + \phi_n).$$

$$\phi_n = \phi_0 + \Delta\phi_n$$



$$\alpha(t) - \alpha(0) = \frac{i}{m\omega_0 a} \int_0^t e^{i\omega_0 t} f(t) dt.$$



Bloch Sphere representation

Experimental realisation: equal rotations applied to each qubit.

Results shown are for experiments performed on an entangled state.



(Our measurement method does not distinguish from $\uparrow\downarrow$ but this has little influence for this example)